

The XMM-LSS survey: The XMDS/VVDS 4σ catalogue ^{*}

L.Chiappetti¹, M.Tajer^{2,5}, G.Trinchieri², D.Maccagni¹, L.Maraschi², L.Paioro¹, M.Pierre³, J.Surdej⁴, O.Garcet⁴, E.Gosset⁴, O.Le Fèvre⁶, E.Bertin⁷, H.J.McCracken⁷, Y.Mellier⁷, S.Foucaud¹, M.Radovich⁸, V.Ripepi⁸, and M.Arnaboldi⁹

¹ INAF IASF, Sezione di Milano “G.Occhialini”, via Bassini 15, I-20133 Milano, Italy
e-mail: lucio@mi.iasf.cnr.it

² INAF Osservatorio di Brera, via Brera 28, I-20121 Milano, Italy

³ CEA/DSM/DAPNIA Service d’Astrophysique, Saclay, F-91191 Gif sur Yvette, France

⁴ Institut d’Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août 17, B-4000 Liège 1, Belgium

⁵ Università degli Studi di Milano - Bicocca, Dipartimento di Fisica, Piazza della Scienza 3, I-20126 Milano, Italy

⁶ Laboratoire d’Astrophysique de Marseille, Traverse du Siphon, F-13376 Marseille, France

⁷ Institut d’Astrophysique de Paris, 89bis bvd Arago, F-75014 Paris, France

⁸ INAF Osservatorio di Capodimonte, via Moiariello 16, I-80131 Napoli, Italy

⁹ INAF Osservatorio Astronomico di Torino, via Osservatorio 20, I-10025 Pino Torinese, Italy

Received jdate_i / accepted jdate_i

Abstract. We present a first catalogue of X-ray sources resulting from the central area of the XMM-LSS (Large Scale Structure survey). We describe the reduction procedures and the database tools we developed and used to derive a well defined catalogue of X-ray sources. The present catalogue is limited to a sub-sample of 286 sources detected at 4σ in the 1 deg^2 area covered by the photometric VVDS (VIRMOS VLT Deep Survey), which allows us to provide optical and radio identifications. We also discuss the X-ray properties of a larger X-ray sample of 536 sources detected at $> 4\sigma$ in the full 3 deg^2 area of the XMM Medium Deep Survey (XMDS) independently of the optical identification. We also derive the logN–logS relationship for a sample of more than one thousand sources that we discuss in the context of other surveys at similar fluxes.

Key words. X-ray – Surveys – AGN – Catalogues

1. Introduction

The XMDS (*XMM* Medium Deep Survey) is an X-ray survey based on the pooling of a significant fraction of the *XMM-Newton* guaranteed time of three “hardware institutes” (IASF Milano for XMM-EPIC, Liège for XMM-OM and CEA Saclay for XMM-SSC).

The XMDS pointings lie at the heart of the full, larger *XMM* Large Scale Structure (XMM-LSS) Survey (see Pierre et al., 2004), to which we refer for a discussion of the motivations and the choice of the sky field.

About two thirds of the XMDS area are covered in the optical band by the VVDS (VIRMOS VLT Deep Survey) both by UBVRI photometry (Le Fèvre et al., 2004a) and by multi-object spectroscopy with VIMOS (Le

Fèvre et al., 2004b), by an associated radio survey at 1.4 GHz (Bondi et al., 2003) and by a GALEX Early Release Observation (Arnouts et al., 2005).

The XMM-LSS area has been covered by an associated radio survey at 74 and 325 MHz (Cohen et al., 2003) and will be covered by surveys in other bands like SWIRE (Lonsdale et al., 2003), the CFHTLS (see web site¹) and by our own follow up observations (Pierre et al., 2004).

Thus, for these areas, multiwavelength information exists or will be gathered in the future, allowing investigations of large scale structures and source populations (clusters, galaxies and AGN) at medium redshifts.

For internal usage within the XMDS and XMM-LSS consortia, and in the future also for public access, a database has been designed, which is presented in this paper together with the first results obtained from the analysis of the XMDS data.

^{*} The full catalogue illustrated in Table 3 is only available in electronic form at the CDS via anonymous ftp to [cdsarc.u-strasbg.fr](ftp://cdsarc.u-strasbg.fr) (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>

¹ <http://www.cfht.hawaii.edu/Science/CFHLS/>

The layout of the paper is as follows. The database tool devised to store the results of the XMM-LSS and XMDS and associated surveys is described in Section 2. Section 3 contains the analysis procedure (closely following that of Baldi et al., 2002) used to generate source lists from the 19 XMM pointings of the XMDS using a threshold that ensures a small number of spurious sources. Different selection criteria are then applied to obtain samples used for different purposes. A relatively low probability threshold is used to compute the logN–logS relation presented and discussed in Section 4. The analysis of the X-ray characteristics is restricted to sources with a signal to noise ratio larger than 4, and the results are presented in Section 5. Only a fraction of the XMDS area is covered by the VVDS survey. For the latter area the identification procedure based on the photometric VVDS data and the derived optical vs X-ray properties are described in Section 6.

Preliminary accounts of this work have been presented at recent conferences (Maraschi et al., 2003; Tajer et al., 2004).

2. The database and catalogue

2.1. The LSS database

The web site of the XMM-LSS Master Catalogue² has been designed as a front end to access the database containing the catalogues, both as a working tool for the XMDS and XMM-LSS consortia and, in the near future, for public access.

The query interface uses a Java servlet (communicating via JDBC protocol to the underlying MySQL³ database) to manage the permissions of different groups of people to access different (parts of) database tables, to perform selections into multiple tables in an user-friendly way, to have a quick look at the data and to export results as ASCII or FITS files.

The database presently includes *database tables* for the results of the X-ray pipelines, VVDS photometry for objects within a $40''$ box around an X-ray source, radio and other catalogues as reported in Section 6.1; it also provides links to *data products* like X-ray images and exposure maps, finding charts in the I band, radio maps, etc.

The database includes also pre-calculated correlation tables, which link the sequence identifiers of objects in an X-ray table with their neighbours in other tables according to predefined proximity criteria (e.g. a $40''$ box or a $6''$ radius), and greatly speed up queries which involve a couple of tables.

Correlations between more than two tables are managed similarly by multi-column correlation tables generated by our identification procedure described below in section 6.1. They are also used to build *virtual tables*, i.e.

² <http://cosmos.mi.iasf.cnr.it/~lssadmin/Website/LSS>

³ MySQL is an open source database server developed by MySQL AB (see <http://www.mysql.com>)

simultaneous views of columns taken from different individual tables, or even of results of algebraic operations between different columns.

Usage of appropriate MySQL syntax during the composition of a query allows the production of results of arbitrary complexity as output from our database, like e.g. ds9 region files, or the L^AT_EX code for the catalogue table reported here (Table 3).

2.2. The catalogue

In this paper we publish, as a first part of the XMDS catalogue, the results of the identification procedure described in section 6.1, i.e. a catalogue of 286 X-ray sources detected at more than 4σ and located in the VVDS area. Further releases will also include sources already present in our database and detected at a lower S/N and/or located outside the VVDS area.

Our current choice allows us to present a catalogue inclusive of reliable (though not spectroscopically confirmed) identifications and of the basic optical and radio characteristics of the X-ray sources.

The catalogue published electronically at CDS along with the present paper (and available as Online material) includes a rather reduced, manageable number of columns. A printout sample of the catalogue in tabular form is presented in Table 3. Comments on individual sources are presented in Appendix A.

Since our database contains much more information than what is included in the published catalogue, we plan to open access to our database query engine, in accordance with the "legacy policy" of the XMM-LSS survey.

Usage of our database interface will allow the user to make selections on the datasets, to access additional columns (including e.g. X-ray fluxes in all bands, photometry in additional bands, radio fluxes, identification ranks and flags, technical information, etc.) which is not practical to include here, and to generate (and operate upon) expressions involving columns.

The query interface will be accessible from the time the paper is accepted at <http://cosmos.mi.iasf.cnr.it/~lssadmin/Website/LSS/Query> logging in as user `xmids`, password `guest` in workspace `public`.

3. Observations and data reduction

3.1. Observations

The XMDS observations consist of 19 overlapping pointings (typical duration in the range 20–25 ks), covering a contiguous area of about 3 deg^2 . The sky location of the combined field of view (FOV) of the pointings is shown in Fig. 1.

The observations were performed by XMM-Newton (Jansen et al., 2001), with the EPIC MOS (Turner et al., 2001) and pn (Strüder et al., 2001) cameras, between July 2001 and January 2003. A basic journal of observations is reported in Table 1, containing the ESA dataset identifier




Fig. 1. Location of the XMDS pointings on the sky : the approximate FOV footprint (with the pn protruding at the actual roll angle) is shown (solid lines); the gray colour for field G12 indicates that it has not been used (see text). The dashed rectangle indicates the area of VVDS photometry.

that can be used to inquire for further details at the ESA *XMM* archive.

The XMDS fields are surrounded at present by 32 additional shorter (typically 10 ks) pointings (B01 to B32) performed in AO1 and AO2 as part of the XMM-LSS program. More detailed information is available on an ancillary page of the XMM-LSS Master Catalogue web site⁴.

3.2. Generation of the X-ray source list

The reduction of the XMDS data has been performed in Milan using a streamlined version of the pipeline developed by Baldi et al. (2002) for the HELLAS2XMM survey. We refer to the latter paper for the formalism and procedure, and only mention here the main points and the few differences.

We used the *XMM-Newton* Science Analysis System (XMM-SAS) v5.4.1, whose tasks `emproc` and `epproc` now provide a reliable identification of the location of bad pixels. To remove data contaminated by soft proton flares, we used *fixed* thresholds (on the global background above 10 keV) of 0.15 counts s⁻¹ for MOS and 0.35 counts s⁻¹ for pn : as a result of this choice, field G12 could not be analysed because the background was almost constantly above these thresholds. A mosaic of our X-ray images is shown in Fig. 2.

As in Baldi et al. (2002), we ran the source detection (inclusive of background computation) and characteriza-

tion procedure on merged MOS and pn data in *five* different energy bands, which we designate as : A (0.3 – 0.5 keV), B (0.5 – 2 keV), C (2 – 4.5 keV), D (4.5 – 10 keV) and CD (2 – 10 keV). We note that sources detected in the CD band do not result from the simple sum of detections in the separate C and D bands, but from the detection algorithm run on the full 2 – 10 keV band, while quantities referring to the total band ABCD (0.3 – 10 keV) are obtained by summation of the results in the individual bands.

We have used the XMM-SAS tasks `ebxdetect` and `emldetect` to obtain a list of candidate source positions, but net counts and associated parameters are obtained from our own program written according to Baldi et al. (2002). In particular fluxes were computed for a power law spectrum with $\Gamma=1.7$ and $N_H=2.61 \times 10^{20}$ cm⁻² (i.e. the average galactic column density in the XMDS field direction (Dickey & Lockman, 1990)), using conversion factors calculated from response matrices generated consistently with the event selections used in our pipeline. In particular, as in Baldi et al. (2002), we used a conservative pattern selection of single and double events for MOS, and only single events for pn. Since the count rates are derived in regions including a fixed percentage (68%) of the expected flux from a point source (whose radius depends on off-axis positions), our fluxes are correct for point-sources. The fact that a source may fall close to (or on) an inter-CCD gap is taken into account in flux computation by proper usage of the EPIC calibration files, and by the usage of exposure maps in the measurement of the count rate. This should be an acceptable approximation (the same used by Baldi et al., 2002). Anyhow, we record in our database a flag indicating whether a source is close to a FOV edge or to CCD borders or gaps.

For insertion in our database, we retain only sources that have a chance detection probability less than 2×10^{-4} in at least one energy band. In the 19 useful pointings of the survey, we detected 1322 X-ray sources (including multiple detections in overlapping regions, see Table 2), satisfying the probability threshold $P < 2 \times 10^{-4}$. In this work we present subsets chosen with more stringent criteria: (1) all sources detected with chance probability less than 2×10^{-5} to derive the logN–logS relationship (section 4); (2) X-ray sources detected with a signal to noise ratio larger than 4 in any one of the energy bands defined above (where the error σ_S on the net number of counts is calculated using the Poissonian approximation from the gross number of counts cts_{img}

$$\sigma_S = 1 + \sqrt{cts_{img} + 0.75}$$

according to Gehrels (1986) as in Baldi et al. (2002)) to study the mean X-ray properties of the brighter end of the sample (the 4 σ sample, section 5); (3) all sources in the 4 σ sample covered by VVDS photometry, for which we present the catalogue (section 2.2) and we discuss the X-ray/optical properties (section 6).

The numbers of sources in each sample are reported in Table 2.

⁴ <http://cosmos.mi.iasf.cnr.it/~lssadmin/Website/LSS/Anc/goepic.html>

Table 1. Journal of XMDS observations^e

Field	ESA obs id	date	pointing RA	DEC	max exposure ^c (ks)	shift ^d Δ RA (")	Δ DEC (")
G01	011268 0101	28 Jan 2002	02:27:25.4	-04:11:06.4	24.9	-1.60	0.00
G02	011268 0201	14 Jul 2002	02:25:54.2	-04:09:05.6	9.8	0.00	-0.53
G03	011268 0301	19 Jan 2003	02:24:45.6	-04:11:00.8	21.8	-0.53	1.07
G04	010952 0101	29 Jan 2002	02:23:25.3	-04:11:07.6	25.8	-1.60	0.00
G05	011268 0401	02 Feb 2002	02:28:05.1	-04:31:08.1	23.7	0.00	0.00
G06	011268 1301	26 Jul 2002	02:26:34.4	-04:29:00.8	13.0	0.00	0.53
G07	011268 1001	30 Jan 2002	02:25:25.3	-04:31:07.1	23.5	-0.53	-0.53
G08	011268 0501	25 Jul 2002	02:23:54.6	-04:29:00.1	18.3	0.00	0.00
G09	010952 0601	31 Jan 2002	02:22:45.2	-04:31:11.1	22.4	n/a	n/a
G10	010952 0201	29 Jan 2002	02:27:25.4	-04:51:04.4	24.8	-0.54	0.53
G11	010952 0301	02 Feb 2002	02:26:05.1	-04:51:06.1	21.9	0.00	0.53
G12 ^a	010952 0401	01 Feb 2002	02:24:45.4	-04:51:11.2	n/a	n/a	n/a
G13	010952 0501	03 Jul 2001	02:23:13.1	-04:49:03.1	23.9	-2.67	-2.13
G14	011268 0801	31 Jan 2002	02:22:04.1	-04:51:09.7	13.6	n/a	n/a
G15	011111 0101	06 Jul 2001	02:27:54.1	-05:09:02.3	21.2	n/a	n/a
G16 ^b	011111 0201	06 Jul 2001	02:26:34.2	-05:09:03.1	3.9	n/a	n/a
G16 ^b	011111 0701	14 Aug 2001	02:26:35.2	-05:08:46.6	23.7	n/a	n/a
G17	011111 0301	03 Jul 2001	02:25:14.3	-05:09:08.4	22.5	n/a	n/a
G18	011111 0401	03 Jul 2001	02:23:54.1	-05:09:09.7	28.1	n/a	n/a
G19	011111 0501	04 Jul 2001	02:22:34.0	-05:09:02.1	23.7	n/a	n/a

a pointing G12 has not been used because of the high background level (see text)

b pointing G16 has been repeated since the first instance was curtailed

c the reported max exposure is the highest value in the exposure maps of the individual cameras, i.e. net of soft proton flares

d the RA and DEC shifts are the astrometric correction applied (see text) for the fields which overlap the VVDS area and for which optical identification has been performed

e more details available through our ancillary web page <http://cosmos.mi.iasf.cnr.it/~lssadmin/Website/LSS/Anc/xmdsep.html>

Sample	N_{tot}	N_B	N_{CD}
$P < 2 \times 10^{-4}$	1322	1166	419
$P < 2 \times 10^{-5}$	1129	1028	328
4σ (detections)	612	591	158
4σ (independent)	536	518	143
4σ VVDS	286	278	73

Table 2. Total number of sources and sources detected in the B and CD bands respectively for the samples presented in the paper. For the $P < 2 \times 10^{-4}$, $P < 2 \times 10^{-5}$ and the first line of the 4σ sample, numbers refer to detections (i.e. include multiple detections of the same source in different fields), while for the second line of the 4σ sample and for the 4σ VVDS sample numbers refer to *independent* sources.

Note that in the present version we have analysed each pointing independently, in spite of the fact that the FOVs of adjacent pointings overlap. The same object could therefore be detected in more than one field. This fact is recognized only *a posteriori* (see section 5). In what follows we will always specify whether the sample we consider includes multiple detections of the same source.

An independent pipeline is being developed in Saclay for the analysis of the entire XMM-LSS, using a wavelet technique and therefore best suited for extended sources

(Pacaud et al. (2005), see a preliminary account in Pierre et al. (2004)) and will be used as a basis for the future complete XMM-LSS catalogue.

4. The logN–logS relationship

We computed the logN–logS distributions in the 0.5 – 2 and 2 – 10 keV band as follows.

We have considered all sources with a detection probability in the 0.5 – 2 and 2 – 10 keV bands below $P = 2 \times 10^{-5}$. This is well below our acceptance threshold of 2×10^{-4} , ensuring very few spurious sources in the sample while giving us a large dataset to compute the logN–logS.

We have generated a flux limit map for each X-ray field and energy band, which contains the faintest flux at which a source can be detected at the assumed level of significance above the local background. From these we have computed the sky coverage plotted in Fig. 3. This considers all fields as independent areas, consistently with our detection procedure (see section 3.2), so the full sky coverage is the sum of all areas surveyed.

Our faintest flux is $\sim 10^{-15}$ erg cm⁻² s⁻¹ in the 0.5 – 2 keV band and $\sim 7 \times 10^{-15}$ erg cm⁻² s⁻¹ in the 2 – 10 keV band.




Fig. 2. Mosaic of our X-ray images in the total (0.3 – 10 keV) band. No exposure map correction has been applied to this figure.

In Fig. 4 we plot the cumulative logN–logS in the 0.5–2 keV band (upper left panel, 1028 sources) and in the 2–10 keV band (lower left panel, 328 sources). The plots relative to individual fields have also been produced and inspected and do not show significant deviations from the combined distribution.

The shapes of the (cumulative) logN–logS are clearly curved and cannot be represented by a single power law. A fit to the cumulative distribution does not allow a meaningful computation of confidence limits on the parameters, because points and errors are not independent (Murdoch et al., 1973; Crawford et al., 1970). Therefore we computed the differential logN–logS by binning (in logarithmic space) the number $N(F)$ of sources with flux F into flux bins of width ΔF_i (i.e. between fluxes F_{jmin} and F_{jmax}), then computing the average sky coverage A_i in the bin. The i -th bin of the differential curve $n(F)$ (shown in the right panels of Fig. 4) is given by

$$n(F)_i = \sum_{F_{jmin}}^{F_{jmax}} \frac{N(F_j)}{\Delta F_i A_i}$$

A statistical error due to the Poissonian error on the number of sources is assigned to each bin. Again, we confirm

that a single power law does not give a good fit. Therefore we used a broken power law.

For the 0.5 – 2 keV band, the best fit is

$$n(F) = \begin{cases} 6.515 \times 10^3 \times F_{15}^{-2.62} & F_{15} > 10.58 \\ 384.2 \times F_{15}^{-1.42} & F_{15} < 10.58 \end{cases}$$

All fluxes F_{15} are normalized to $10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$. 90% confidence limits on slopes are $2.62^{+0.25}_{-0.22}$ and $1.42^{+0.14}_{-0.15}$, while the break position is in the range $1.06^{+0.30}_{-0.22} \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$.

For the 2 – 10 keV band we find

$$n(F) = 4.483 \times 10^4 \times F_{15}^{-2.91}$$

for $F_{15} > 21.4$, with 90% errors on slope $2.91^{+0.45}_{-0.30}$ and break at $2.14^{+0.81}_{-0.54} \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$. At lower fluxes the slope is not well constrained: the best fit is 1.53, but the confidence interval ranges between 0.37 and 2.04.

In Fig. 4 we show the fit results both in the differential and the cumulative forms for a better comparison with the literature that gives sometimes cumulative (e.g. Baldi et al., 2002; Moretti et al., 2003), sometimes differential plots (e.g. Yang et al., 2004; Harrison et al., 2003). Given the uncertainties in the fits, we prefer the comparison with the actual data where possible.




Fig. 4. LogN-logS of sources detected at $P < 2 \times 10^{-5}$ in band B in the cumulative (upper left panel) and differential (upper right panel) form and the same in band CD (lower left and right panels). The units of $n(F)$ are number per $10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ and deg^{-2} . Our best fits are reported as solid lines on the differential plots and have also been converted to the cumulative form.

Our probability threshold $P < 2 \times 10^{-5}$ is one of those used in Baldi et al. (2002). Since our procedure is very similar to theirs and the flux range is nearly the same, the logN-logS should in principle be almost identical. We find consistency (within errors) with the integral HELLAS2XMM distributions for the lower fluxes: $F_{0.5-2} \lesssim 3 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ and $F_{2-10} \lesssim 2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$, while at higher fluxes we are nominally lower, but, taking into account the purely statistical error on the number of sources, we are consistent with the lower envelope of the Baldi et al. (2002) relationship.

A good agreement is also found with the *Chandra* Large Synoptic X-ray Survey (CLASXS, Yang et al., 2004) and with the Serendipitous Extragalactic X-ray Source Identification (SEXSI, Harrison et al., 2003) program, in the common flux range. We compared the differential plots and found consistency with the CLASXS survey in the soft band (no plot in that band is given in Harrison et al., 2003). In the hard band we are consistent with the SEXSI survey, but we are systematically lower than the CLASXS points; this is not surprising, given that Yang et al. (2004) find that their total counts at $F_{2-8} \sim 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ are $\sim 70\%$ higher than those of Moretti et al.




Fig. 3. Sky coverage for sources detected at $P < 2 \times 10^{-5}$ in the B band (solid line) and in the CD band (dashed line) computed considering all fields as independent (see text).

(2003), who use Baldi et al. (2002) data for the flux range we are interested in.

The sky location of the XMDS survey has been intentionally chosen in order to avoid known bright X-ray sources and this could therefore explain, at least partially, the low density observed in our survey for fluxes above a few 10^{-14} erg cm $^{-2}$ s $^{-1}$. On the other hand, Yang et al. (2004) interpret their overdensity as an indication of an underlying large scale structure. We note that the CLASXS, like the XMDS, is constructed on contiguous pointings, while the HELLAS2XMM and the SEXSI surveys are serendipitous surveys: it seems reasonable that the number count difference we find between the different surveys is caused by cosmic variance.

5. X-ray properties of the total XMDS 4σ sample

One of our criteria for the definition of a sample is the presence of a localized X-ray count excess with signal to noise ratio (as defined in section 3.2) ≥ 4 in at least one of the energy bands (the “ 4σ sample”). This results in 612 *detections*. Taking into account multiple detections (same source detected in different overlapping fields) the total number of *independent sources* reduces to 536. All detections in different pointings with X-ray positions closer than $6''$ have been automatically considered as multiple detections of the same source. For larger separations (up to $18''$, which can be considered as a boundary for suspicious overlaps, since there are only 2 pairs of sources detected in the *same* field which are closer than such a distance, one at $8.7''$ and one at $15.9''$), we examined each of the 8 pairs and recovered 2 cases: one is included in our catalogue and is discussed in Appendix A, the other

has a distance of $6.4''$, but one detection is outside the FOV of one camera. For all multiple detections we used the position and derived parameters from the detection that gave the highest significance, unless the source falls outside the FOV of one camera, or close to an inter-CCD gap. We do not detect any significant flux variation in the multiply detected sources with the possible exception of three, which are however close to inter-CCD gaps. The apparent flux decrease in the $0.5 - 2$ keV band from the brightest observation in each couple would be of -35% , -65% and -28% , with a significance of ~ 3.5 , 3.9 and 4.5 σ respectively.

5.1. Fluxes and hardness ratios

The flux distribution of the sources in the 4σ sample is shown in Fig. 5 for sources detected in the energy bands $0.5 - 2$ (left panel) and $2 - 10$ keV (right panel) respectively.

We find that 518 sources ($\sim 97\%$) are included in the sample because of their detection in the $0.5 - 2$ keV (B) band, while only 143 ($\sim 27\%$) satisfy the detection criterion in the $2 - 10$ keV (CD) band. Clearly most of the latter are *also* detected in the B band. We do not have any source included in the sample because of its detection in the $4.5 - 10$ keV band *only*, and only 16 hard sources ($\sim 10\%$) are not detected in the $0.5 - 2$ keV band. As indicated by Fig. 5, we find *a posteriori* $F_X \sim 2.5 \times 10^{-15}$ erg cm $^{-2}$ s $^{-1}$ as the flux limit in the B band and $F_X \sim 2 \times 10^{-14}$ erg cm $^{-2}$ s $^{-1}$ in the CD band.

A spectral analysis of the brighter sources is in progress. Here we present the results of the hardness ratio analysis that can be done for a significantly larger number of sources, and is the only spectral analysis possible for the fainter sources in the sample.

To this end we note that, even if a source does not satisfy the detection threshold in a band different from the one that motivated the inclusion in the sample, we have an estimate of the counts and fluxes in all bands considered, which we can use to get spectral information. Let us define:

$$HR = \frac{CR_{2-10} - CR_{0.5-2}}{CR_{2-10} + CR_{0.5-2}}$$

$$HR_{cb} = \frac{CR_{2-4.5} - CR_{0.5-2}}{CR_{2-4.5} + CR_{0.5-2}}$$

$$HR_{dc} = \frac{CR_{4.5-10} - CR_{2-4.5}}{CR_{4.5-10} + CR_{2-4.5}}$$

where CR are the net count rates in the given energy band corrected for PSF and vignetting. We only include sources for which we have a reasonable estimate of the flux (i.e. a signal-to-noise ratio ≥ 2) in the $4.5 - 10$ keV band: upper limits in such a band do not carry significant information since they provide only an upper limit to HR_{dc} , which is expected for fainter sources with the canonical unabsorbed power law spectrum. On the contrary upper limits from




Fig. 5. Flux distribution in band B (left panel) and in band CD (right panel) for the total 4σ sample. The hatched area gives the distribution of the sources detected at $\geq 4\sigma$ in the concerned band.




Fig. 6. Hardness ratio HR vs 2 – 10 keV flux for all sources detected at $\geq 4\sigma$ in the CD band (asterisks). Only a few typical error bars are shown to avoid figure overcrowding, inclusive of all those detected at $\leq 2\sigma$ in the B band. Dashed lines indicate hardness ratios expected for an absorbed power law spectrum with energy index $\Gamma = 1.7$ and intrinsic N_H as reported in the label.

the 0.5 – 2 keV band are interesting as they provide lower limits to HR_{cb} .

We verified that the values obtained from the combined count rates from different cameras are consistent with those computed from single camera data.

In Fig. 6 we plot the main hardness ratio HR as a function of the 2 – 10 keV flux. Also shown in the figure are dashed lines indicating the hardness ratios expected for a simple power law model with spectral index $\Gamma = 1.7$ and increasing values of the absorbing column N_H . This plot shows that $\sim 30\%$ of sources detected in the CD band have hardness ratios consistent with $N_H > 10^{21.5} \text{ cm}^{-2}$, for $z = 0$. Note that if sources are at higher redshifts, the horizontal lines for different values of N_H shift downwards, i.e. the same hardness ratio corresponds to a higher absorption column.

Within our current sample, we do not find a spectral hardening with decreasing X-ray flux as reported by other authors; however our statistics in the hard band is lower than those from other surveys. On the other hand, it appears from numbers in Table 2 that the fraction of sources in the hard band increases when the sample is larger (from 26% in the 4σ sample to 32% in the total $P < 2 \times 10^{-4}$ sample). Since a less restrictive threshold implies a lower flux limit, we can infer that the fraction of hard sources should increase as the flux decreases.

Fig. 7 shows the so-called X-ray colour-colour diagram, i.e. the plot of the hardness ratio between the harder bands HR_{dc} vs the hardness ratio between soft and hard bands HR_{cb} .

Sources with HR consistent with $N_H > 10^{22} \text{ cm}^{-2}$ in Fig. 6 occupy the same locus in Fig. 7 (or $N_H > 10^{23} \text{ cm}^{-2}$ if we consider that sources are at $z \sim 1$). This singles them out as promising candidates for more detailed studies.

In spite of the relatively small number of objects, it is worth noting that a few sources with low values of HR_{cb} and high values of HR_{dc} are present: these are not consistent with a simple power law model. In particular, their




Fig. 7. Hardness ratio HR_{dc} vs HR_{cb} for sources in the $\geq 4\sigma$ sample (left panel), which are also above 2σ in band D. Typical error bars are shown (dashed when the significance in the involved bands is below 4σ). Grids in right panel represent hardness ratios expected with a simple power law model for values of spectral index and N_H reported in the labels at $z = 0$ (empty circles and dashed lines) and at $z = 1$ (full squares and dotted lines).

location in the diagram at $HR_{dc} > -0.2$ and $HR_{cb} < -0.5$ could suggest a concave spectral shape, due to the relative lack of signal around 2 keV. Such a shape could result from the superposition of an unabsorbed component on a flat/absorbed power law as often observed in obscured AGNs, where the unabsorbed component can be due to scattered or transmitted AGN continuum or to thermal and non thermal emission associated with starburst regions.

6. The XMDS/VVDS 4 σ sample

6.1. The identification procedure

For the generation of our sample of candidate identified X-ray sources we have used the following data sources (database tables and data products).

- The database table of X-ray sources from our pipeline.
- The optical tables with the VVDS UBVRI(JK) photometry including both sources with reliable photometric data and objects whose photometry is flagged as unreliable because the objects are saturated, masked or close to the borders of the VVDS area. In addition, we have used the associated finding charts. Data quality and limiting magnitudes are discussed in McCracken et al. (2003) and Radovich et al. (2004).
- The table with the VIRMOS 1.4GHz radio catalogue (Bondi et al., 2003) which covers the 1 deg² VVDS area at a depth of 80 μ Jy and a resolution of 6".
- The table with the XMM-LSS radio catalogue (Cohen et al., 2003), a shallow survey at 74 (4 mJy, 5.6 deg²) and 325 MHz (275 mJy, 110 deg²) over a larger area

encompassing the XMM-LSS fields. Occasionally we used the associated radio maps.

- The table with the tentative cluster candidates
- The NED and SIMBAD database tables, containing pointers to objects within 20" from an X-ray source. It should be noted that the VIRMOS 1.4GHz catalogue has recently been inserted also in SIMBAD, which could result in a redundant association.

The identification process has followed a multi-step procedure, that also involved the determination of reliable X-ray/optical angular separation criteria for identifications and the testing of more automated identification procedures for future works.

First of all, we used our database to create the sample of X-ray sources that are within the VVDS area (i.e. for which we can provide a FITS finding chart, covering a 40" box around the X-ray position, that we visually inspected).

For each unique X-ray source we have then conservatively considered all optical objects within a radius of 6" from the X-ray position (see e.g. Waskett et al., 2004). This radius is considerably larger than the nominal SAS (`emldetect`) position error combined with the 2" XMM astrometrical uncertainty. For such objects we have positions, magnitudes and a crude optical classification (star-like vs extended) from the VVDS database.

We have identified the optical counterparts according to the following criteria: a) unique optical counterpart [67 objects]; b) closest and brightest ($\Delta\text{mag} > 2$) counterpart [166 objects]; c) no optical counterpart down to $I \sim 25$, the magnitude limit of McCracken et al. (2003) [5 objects, of




Fig. 8. Examples from our identification procedure : some representative I band finding charts (size 40'' $\times$ 40'') with overlays of the X-ray error circles (white thick at 4'' radius), the positions of all optical sources within 6'' (cyan X's and crosses respectively for point-like and possibly extended objects), the positions of NED catalogue objects (yellow diamonds) and the radio source error box (green rectangles) or extent (green ellipses). The printed figure is in grayscale : see the online edition for a colour version. Three cases are shown : an unique counterpart, an identification preferring a radio source, an ambiguous identification.

which 3 with no counterpart at all, 2 with a radio counterpart]. We also used the presence of a radio counterpart as a preference criterion. The remaining cases [55] do not allow us to assign an unambiguous optical counterpart until spectroscopic confirmation of the source is available, so all possible optical counterparts are retained. 22 cases were flagged as suspect groups, clusters or simply crowded fields, of which half have however a dominant, unique counterpart. Examples of finding charts used in our identification procedure are shown in Fig. 8.

After a first pass with this procedure, we found evidence in some X-ray fields of a systematic rigid shift between the position of the X-ray sources and their counterparts (as high as 3'' in field G13). We profitably used the pairs identified at such stage to compute (via the SAS task *eposcorr*) a rigid shift correction (see Table 1), and inserted in the database two additional columns with corrected X-ray positions. No rotational correction was required. See also section 6.2 and Fig. 9. After this astrometric correction we repeated our inspection and ranking procedure.

This time-consuming identification process involved a number of iterations and intermediate steps, which we have recorded in the database with different flags and fields (that could identify for example the presence of a visually crowded field, the cross-correlation with a detection of the same source in a different field, the source and quality of the optical photometry). These will remain as part of the record for each source for future automatic database queries, but are not included in the catalogue attached to this paper.

The final identifications are reported in the catalogue presented here and have associated numerical codes that describe the quality (rank) of the identification, as described in point M of Table 3.

As a final step, we have computed the probability of chance coincidence between the X-ray source and the optical candidates within the given radius according to the

formula by Downes et al. (1986) used e.g. by Brusa et al. (2003) or Mignoli et al. (2004)

$$p = 1 - \exp(-\pi n(< m) r^2)$$

where r is the distance between the X-ray source and the concerned optical candidate, and the density $n(< m)$ of optical objects having I magnitude brighter than the magnitude m of the candidate counterpart was computed using the full VVDS catalogue (McCracken et al., 2003). A $p < 0.01$ criterion (similar to the one used by Brusa et al. (2003) and more conservative than the one used by Mignoli et al. (2004), corresponding in our case e.g. to the probability of finding an I=21 object at 2'') applied to the X-ray/optical pairs produces a list of identifications very similar to our visually identified sources. Namely 93% of our best identifications (rank 0) and 84% of the rank 1 associations have $p < 0.01$ (and only one of the rank 0 identification has $p \geq 0.02$), while only 21% of the ambiguous (rank 2) cases have $p < 0.01$.

At the present time, spectroscopy is in progress within the framework of different programs, and only a few of our identifications are spectroscopically confirmed.

6.2. Optical vs X-ray properties

Fig. 9 shows the angular separation between the position of the best optical counterpart identified according to the positional and photometric criteria (as described in section 6.1) and the position of the X-ray source (after astrometric correction).

One can note the absence of systematic shifts. We also verified that there is no correlation between the X-ray to optical distance and the off-axis position in the field of view, while there is a mild correlation with count rate, in the sense that sources with a large X-ray to optical separation are among the weakest. We note moreover that 93% of the optical counterparts are within a 2'' radius from the X-ray source (99% within 4''). We therefore conclude that,




Fig. 9. Distance in RA and DEC between the X-ray source position (inclusive of astrometric correction) and the optical counterpart for well identified sources (ranks 0 and 1 in the database). Two fiducial radii of 2'' and 4'' are shown.




Fig. 10. R magnitude distribution for well identified sources (ranks 0 and 1 in the database, solid histogram) and for optical objects within 6'' from the ambiguously identified sources (rank 2, dashed histogram)

after astrometric corrections have been applied, the search radius for optical counterparts can be safely reduced to 4''.

Figure 10 compares the distribution of optical magnitudes for the identified sources and for the ambiguous identifications, i.e. X-ray sources for which there are 2 or more candidate identifications in the search radius.




Fig. 11. X-ray to optical flux ratio (band CD vs R magnitude) as a function of the X-ray flux (in band CD). The sources shown are those in the 4 σ VVDS sample, detected above 2 σ in band CD. Upper limits (on the X-ray to optical ratio) correspond to sources photometrically saturated (optical magnitude likely underestimated). Lower limits corresponds to sources without optical counterpart (1 association with a radio source only). Open symbols correspond to rank 2 sources (ambiguous identification, more than a counterpart, of which the brightest is used to derive the optical flux). Diagonal lines indicate loci of constant R magnitude while horizontal dotted lines mark the region of canonical AGNs ($0.1 < F_X/F_R < 10$)

As expected, unidentified sources populate the region of fainter magnitudes where the density of possible optical counterparts makes chance coincidences more probable. Spectroscopy or at least multicolor photometry is needed to increase the number of identifications and to determine the redshifts and types of the candidate counterparts. Work on the optical and multifrequency information already acquired within the VIMOS and XMM-LSS consortia is in progress.

We compute the X-ray to optical flux ratio using our R magnitudes (in the AB system) according to the formula

$$\log(F_X/F_R) = \log F_X + 5.51 + R_{AB}/2.5$$

where the constant derives from the standard conversion of AB magnitudes into monochromatic flux and from the integration of the monochromatic flux over the filter bandwidth assuming a flat spectrum.

The X-ray to optical ratio for our identified sources is shown in Fig. 11 as a function of the X-ray flux (2 – 10 keV band). We have also included the ambiguous identifications (marked with different symbols) assuming that the brighter object is the true counterpart. The latter values should therefore be considered as lower limits for the X-ray to optical flux ratio: a few objects that are now




Fig. 12. X-ray (2 – 10 keV) to optical flux ratio vs hardness ratio for sources of the 4 σ VVDS sample. Only sources with a significance of at least 2 σ in one band (and of course 4 σ in the other) are shown, provided they have a rank 0-1 identification (one case has only a radio counterpart). Upper limits are due to “saturated” optical counterparts which only yield a lower limit to the optical flux.

in the “normal AGN” region could migrate upwards and populate the obscured AGN region.

As expected, most ($\sim 80\%$) of the sources lie in the region of classical AGNs ($0.1 < F_X/F_R < 10$), $\sim 3\%$ are in the starburst – normal galaxy range ($F_X/F_R < 0.1$) and $\sim 17\%$ (including optically blank fields) in the region expected for optically obscured AGNs ($F_X/F_R > 10$).

The percentage of high X-ray to optical ratio sources is only slightly lower than in other X-ray surveys of similar depth (e.g. 20% in Fiore et al., 2003). However, since the 4 σ VVDS sample is not fully identified and since our identification procedure favours optically brighter objects, the real fraction of sources with $F_X/F_R > 10$ could be higher.

Fig. 12 shows the X-ray to optical ratio vs the hardness ratio. As shown recently by Della Ceca et al. (2004) this diagram is a useful tool to separate stars from the extragalactic population, and to locate different kinds of AGN. Since we have used the X-ray flux in the 2 – 10 keV band (for comparison with Fig. 11), stars are not included in our diagram as they are not detected in the 2–10 keV band. We verified however that our confirmed stars (via SIMBAD or via preliminary results of VIMOS spectroscopy) fall in the expected location in the $F_{0.5-4.5}/F_R$ vs HR_{cb} plot. Most of our sources populate the expected locus of AGNs, with a number of objects with very high (10-100) X-ray to optical ratios (suggesting optical extinction) and a few with hard X-ray spectra, pointing to X-ray absorbed nuclei. In a forthcoming paper we will take advantage of the full multifrequency and spectroscopic in-

formation to discuss the properties of the populations selected by the X-ray survey with respect to those of the optical survey.

Acknowledgements. This work is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and the USA (NASA).

This work has benefitted from many useful discussions with S. Andreon, A. Wolter and R. Della Ceca at INAF-OABrera.

OG, EG and JS would like to acknowledge support by contract Inter-University Attraction Pole P5/36 (Belgium) and by PRODEX (XMM-LSS)

References

- Arnouts, S., et al. 2005, *ApJ*, 619, L43
- Baldi, A., Molendi, S., Comastri, A., et al. 2002, *ApJ*, 564, 190
- Bondi, M., et al. 2003, *A&A*, 403, 857
- Brusa, M., et al. 2003, *A&A*, 409, 65
- Cohen, A. S., et al. 2003, *ApJ*, 591, 640
- Crawford, D. F., Jauncey, D. L., & Murdoch, H. S. 1970, *ApJ*, 162, 405
- Della Ceca, R., et al. 2004, *A&A*, 428, 383
- Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
- Downes, A. J. B., Peacock, J. A., Savage, A., & Carrie, D. R. 1986, *MNRAS*, 218, 31
- Fiore, F., et al. 2003, *A&A*, 409, 79
- Gehrels, N. 1986, *ApJ*, 303, 336
- Harrison, F. A., Eckart, M. E., Mao, P. H., Helfand, D. J., & Stern, D. 2003, *ApJ*, 596, 944
- Jansen, F., et al. 2001, *A&A*, 365, L1
- Le Fèvre, O., et al. 2004a, *A&A*, 417, 839
- Le Fèvre, O., et al. 2004, *ArXiv Astrophysics e-prints*, astro-ph/0409133
- Lonsdale, C. J., et al. 2003, *PASP*, 115, 897
- Maraschi, L., et al. 2003, *Conference on Multiwavelength AGN Surveys*, Dec 2003, Cozumel
- McCracken, H. J., et al. 2003, *A&A*, 410, 17
- Mignoli, M., et al. 2004, *A&A*, 418, 827
- Moretti, A., Campana, S., Lazzati, D., & Tagliaferri, G. 2003, *ApJ*, 588, 696
- Murdoch, H. S., Crawford, D. F., & Jauncey, D. L. 1973, *ApJ*, 183, 1
- Pacaud, F., et al. 2005, in preparation
- Pierre, M., et al. 2004, *J. Cosmology Astropart. Phys.*, 09, 011
- Radovich, M., et al. 2004, *A&A*, 417, 51
- Strüder, L., et al. 2001, *A&A*, 365, L18
- Tajer, M., Chiappetti, L., Trinchieri, et al. 2004, 6th National Conference on AGNs, May 2004, Volterra, <http://www.arcetri.astro.it/~agn6/proceedings/>
- Turner, M. J. L., et al. 2001, *A&A*, 365, L27
- Waskett, T. J., Eales, S. A., Gear, W. K., et al. 2004, *MNRAS*, 350, 785
- Yang, Y., Mushotzky, R. F., Steffen, A. T., et al. 2004, *AJ*, 128, 1501

Table 3. Sample of our X-ray catalogue. We report here only the first few lines of the catalogue to show the layout. The full catalogue will be available at CDS, as "Online material" or through our database as described in Section 2.2. Comments on peculiar sources are reported in Appendix A.

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	
580	XMDS J022315.4-042558	02 23 15.5 -04 25 58.4	56.7 11.8 22.1 10.5 10.0 20.3	B 1 1 1.4																1	APMUKS (BJ) B022044.68-043934.9	580	
839	XMDS J022315.3-044320	02 23 15.5 -04 43 18.4	48.2 10.8 22.3 9.8 3.9 9.0	B 0 2 2.1 25.45 0.16 24.24 0.36	1																	839	
310	XMDS J022316.3-042007	02 23 16.4 -04 20 07.8	121.8 15.7 1.0 7.2 12.2 0.5	B 0 0 1.8																1	BD-04 388	310	
846	XMDS J022317.2-044035	02 23 17.4 -04 40 33.5	167.2 18.0 72.4 13.3 17.0 37.3	B 0 1 1.1 18.47 0.00 18.23 0.00	1																	846	
828	XMDS J022318.8-044616	02 23 19.0 -04 46 13.9	166.7 17.9 53.0 11.5 11.0 17.4	B 0 0 0.2 21.16 0.01 20.77 0.01	0																	828	
321	XMDS J022319.2-041648	02 23 19.4 -04 16 48.4	55.8 11.4 9.6 8.4 4.1 3.6	B 0 1 1.1 22.62 0.04 21.73 0.03	0																	321	
820	XMDS J022319.4-044732	02 23 19.6 -04 47 30.4	280.6 22.5 125.6 16.0 17.4 38.7	B 0 0 0.7 19.14 0.01 18.49 0.00	0																	820	
779	XMDS J022321.8-045740	02 23 22.0 -04 57 38.4	20.1 9.1 98.7 14.7 1.9 47.2	CD 0 0 1.0 20.35 0.01 19.47 0.01	1																	779	
333	XMDS J022324.9-041452	02 23 25.1 -04 14 52.2	70.5 12.9 28.5 9.8 4.7 9.4	B 0 1 0.6 21.83 0.02 21.31 0.02	0																	333	
550	XMDS J022326.0-043537	02 23 26.0 -04 35 37.1	115.3 15.3 31.3 10.6 17.7 25.1	B 1 1 1.0 23.16 0.04 22.33 0.04	0																	550	
315	XMDS J022326.3-041838	02 23 26.5 -04 18 38.5	51.6 11.5 8.9 8.4 4.6 4.0	B 0 0 0.5 24.29 0.08 23.27 0.07	0																	315	
782	XMDS J022326.3-045708	02 23 26.5 -04 57 05.1	122.8 15.8 62.2 12.3 11.4 29.1	B 0 0 1.5 20.74 0.01 20.49 0.01	1																	782	
789	XMDS J022329.1-045452	02 23 29.3 -04 54 50.8	245.2 21.2 52.0 11.5 22.0 24.0	B 2 1 1.2 20.25 0.01 19.92 0.01	0																	789	
789																							789
571	XMDS J022330.2-043004	02 23 30.3 -04 30 04.2	121.7 15.7 47.0 11.7 14.0 27.6	B 1 0 1.7 20.18 0.01 19.94 0.01	0																	571	
825	XMDS J022330.6-044633	02 23 30.8 -04 46 31.8	303.2 23.3 100.4 14.8 20.5 33.6	B 0 1 0.8 24.44 0.12 23.37 0.08	0																	825	
840	XMDS J022330.9-044235	02 23 31.1 -04 42 33.0	59.1 12.4 37.6 10.6 6.3 20.8	B 2 2 5.4 24.77 0.14 24.59 0.24	0																	840	
780	XMDS J022332.0-045740	02 23 32.2 -04 57 38.7	184.5 18.7 49.5 11.6 18.1 24.6	B 0 0 0.7 19.99 0.01 19.63 0.01	1																	780	
354	XMDS J022332.2-041022	02 23 32.3 -04 10 22.1	128.4 16.2 9.3 7.6 7.7 2.8	B 0 0 0.3 17.77 0.00 16.93 0.00	2																	354	
807	XMDS J022333.0-044924	02 23 33.2 -04 49 22.5	143.8 16.7 48.0 11.3 11.2 17.9	B 2 1 0.9 20.60 0.01 20.26 0.01	0																	807	

A internal sequence identifier

B IAU catalogue name of the object (built from uncorrected coordinates as provided by the XMM-SAS) ; the X-ray properties of a source, i.e. columns from B to L, appear once only on the first catalogue entry for such a source. If the X-ray source has further possible counterparts, these columns are blank and their value is the same as the first entry with same sequence identifier.

C astrometrically corrected RA (J2000)

D astrometrically corrected Declination (J2000)

E number of net counts in band B

F error on net counts in band B

G number of net counts in band CD

H error on net counts in band CD

I flux in 10^{-15} erg cm $^{-2}$ s $^{-1}$ in band B

J flux in 10^{-15} erg cm $^{-2}$ s $^{-1}$ in band CD

K band in which S/N ratio is the highest

L gap flag : can assume a value of 0 (good), 1 (fair), or 2 (poor) related to the fact that the X-ray source falls on inter-CCD gaps in one or more of the EPIC cameras, or close to the edge or outside the FOV of one EPIC camera and is based on the ratio of the punctual exposures in each camera to the maximum exposure in the FOV also for each camera. It can also assume the value -1 if the X-ray source is outside the FOV of at least one camera (there are no such cases in the 4 σ sample).

M identification rank : gives the quality of the identification with the counterpart, coded as 0 optimal e.g. only one counterpart, or brightest and closest; 1 good; 2 (dubious or ambiguous, e.g. typically more than one possible counterpart listed); 4 unidentified source

N distance in arcsec between X-ray source and counterpart in the following order of preference : the VVDS optical counterpart if any; or the radio counterpart in the VIRMOS 1.4GHz catalogue (Bondi et al., 2003); or the radio counterpart in the XMM-LSS 325 MHz catalogue (Cohen et al., 2003); or the catalogued NED object; or the catalogued SIMBAD object. It is blank if there are no counterparts.

O R magnitude of counterpart from the VVDS catalogue. R and I magnitudes are non-blank when there is a valid measurement in the VVDS catalogue. They may be left blank also in case of radio counterparts only, or of (usually very bright) objects present *only* in external (SIMBAD) catalogues.

P error on R magnitude

Q I magnitude of counterpart

R error on I magnitude

S photometry reliability flag (derived from McCracken et al., 2003) : 0 good; 1 flagged as possibly bad, but not saturated; 2 flagged as bad because saturated

T a V indicates presence of a radio counterpart in VIRMOS 1.4GHz catalogue (Bondi et al., 2003)

U an L indicates presence of a radio counterpart in the XMM-LSS 325 MHz catalogue (Cohen et al., 2003)

V name of counterpart, precedence order is NED name, then SIMBAD name including VIRMOS 1.4GHz catalogue then name in the radio catalogue by Cohen et al. (2003)

W position of identification in Appendix A

Appendix A: Comments on individual sources

- XMDS J022315.4–042558: it is identified with a bright spiral, present also in the NED database ($b_J=19.75$), which however falls in a bad photometric area (close to VVDS border) and has no valid optical magnitudes.
- XMDS J022316.3–042007: it has as counterpart the bright ($V=11.0$) star BD–04 388 which also falls in a bad photometric area close to VVDS border, and has no valid optical magnitudes ; even if it didn't, it would have been saturated. The distance reported in the catalogue is overestimated, since the bright source in the VVDS catalogue is split in two by a saturated spike.
- XMDS J022330.9–044235: it is considered a dubious (rank 2) identification, since the closest object reported as rank 2 is faint ($I=24.5$) and at $5.4''$. It is a borderline case, which could perhaps be a blank field.
- XMDS J022337.3–040938: it coincides with a bright saturated ($I < 14.6$) starlike object, which matches a 7.9 mJy 325 MHz radio source, which is a lobe component of the multiple source CRK2003 J0223.6–0409. Another component is instead close to the NED object NVSS J022336–040932. Could it be that the star is a foreground object with respect to the radio source ? The X-ray optical distance reported in the catalogue is overestimated, since the bright source in the VVDS catalogue is split in two by a saturated spike.
- XMDS J022403.7–043303: it has as counterpart the 16th magnitude object Markarian 1036, which is unfortunately saturated in the VVDS.
- XMDS J022403.9–041326: it falls inside a group of weak uncatalogued objects, which are affected by the spike of a bright star situated north of them. A radio object detected both in the VIRMOS 1.4GHz catalogue and in our own observations at 325 MHz (CRK2003 J0224.0–0413) is at $13.6''$ south. This source has a duplicate detection in a different pointing at $11.7''$, and both are inside an unpublished galaxy cluster (Andreon, private communication).
- XMDS J022413.2–045723: it falls at a blank position in a small group of objects, unconfirmed yet as a spectroscopic group, of which the closest is at $2.6''$. Note that this source is in field G18, which is not yet astrometrically corrected.
- XMDS J022445.4–043656: it is considered unidentified (rank 4), falling in a blank field since the closest object (rejected) is faint ($I=25.7$) and at $5.0''$.
- XMDS J022450.9–042903: its brightest and closest counterpart is a 21 mag object at $0.5''$, however at $2.8''$ there is a radio source detected in the VIRMOS 1.4GHz survey as well as in our own 325 MHz data (CRK2003 J0224.8–0429). We kept it as a rank 2 source.
- XMDS J022454.7–040628: its brightest and closest counterpart is a 21 mag object at $1.1''$, however there is a fainter object at $2''$ which has also 1.4 GHz radio emission.
- XMDS J022456.0–041725: it appears to match positionally an uncatalogued enhancement in the periphery of the NED spiral galaxy 2MASX J02245534–0417351 (distance of the nucleus from the X-ray source $14.8''$).
- XMDS J022524.8–044042: it is associated with a saturated bright elliptical, also listed in NED ($b_J=20.04$) and as a VIRMOS 1.4GHz radio source.
- XMDS J022530.6–041418 is one of the sources mentioned in section 5 for which the observations in two different pointings give a marginal flux difference (3.5σ). Its counterpart is a saturated object (nominal VVDS $I=16.21$), corresponding to the NED elliptical galaxy 2MASX J02253077–0414183.
- XMDS J022544.6–041936 is the other source in the VVDS/ 4σ catalogue mentioned in section 5 for which the observations in two different pointings give a marginal flux difference (3.9σ). Its counterpart is also a saturated object (nominal VVDS $I=17.49$), corresponding to the NED elliptical galaxy APMUKS(BJ) B022313.84–043304.3.
- XMDS J022606.6–040259: it is assigned rank 2 because the likely counterpart is not too close to the X-ray position and appears to be an unresolved double object.
- XMDS J022606.9–043316: its counterpart should be a background object masked behind the bright ($V=10.37$) star BD–05 456 (which is too far to be the counterpart). For this reason, the line-of-sight association is classified as rank 2.
- XMDS J022713.6–043910: it has as counterpart the bright ($V=10.12$) star BD–05 461 not included in the VVDS catalogue; if it were, it would have been saturated.
- XMDS J022714.2–042645: the finding chart is saturated by a bright star, BD–05 480, located southwards with respect to our target, whose position is partially behind a spike from the star. For this reason there are no objects listed in the VVDS catalogue, although an object is clearly visible behind the spike at approximately $RA=02:27:14.16$ $DEC=-4:26:45.0$.
- XMDS J022721.7–044153: it has no nearby optical counterparts, but a very good positional coincidence ($0.8''$) with a VIRMOS 1.4GHz radio source. Note that a 17 mag saturated object is however present $6''$ north.
- XMDS J022725.1–043656: it is considered unidentified (rank 4), falling in a blank field since the closest object (rejected) is faint ($I=25.8$) and at $4.2''$.
- XMDS J022735.6–041317: it corresponds to a plethora of very faint objects all located inside an extended radio halo, whose central position matches very well the X-ray position.
- XMDS J022735.7–041122: there are two possible optical counterparts which are both located at the periphery, but inside the nominal extent, of the extended 325 MHz radio source CRK2003 J0227.6–0411 centered $19''$ to the north-east. A 1.4 GHz point-like source, which is a component of a terribly elongated multiple object, is instead located in the X-ray error

radius between the two optical objects, while the NED object NVSS J022736–041117 is located at $9''$ midway between the X-ray source and the 325 MHz radio source.

- XMDS J022742.1–043607: there are no optical catalogued sources; an unresolved double source is present at $4''$ just outside of the nominal error radius of the X-ray source and within $1.1''$ there is a 1.4 GHz radio point source. The finding chart is affected by the spikes of a nearby bright star.
- XMDS J022748.2–041013: it has no optical counterpart at all, hence is flagged as rank 4, blank field.
- XMDS J022812.5–045717: it is in field G15 which has not yet been astrometrically corrected.

Concerning spectral peculiarities, we can use as an indicator the fact that the energy band with the best signal to noise (column K of the catalogue) is not band B. We have only one source (XMDS J022410.7–042759) which is very soft (detected only in band A), and 10 cases with the best signal to noise in band CD (of these 7 are among the 16 not detected in the softer bands mentioned in Section 5.1).

Online Material

We report here the complete file with our catalogue. For an explanation of the columns refer to the sample table 3 in the main text.

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	
580	XMDS	J022315.4-042558	02 23 15.5 -04 25 58.4	56.7 11.8	22.1 10.5	10.0	20.3	B 1 1	1.4												1	APMUKS(BJ) B022044.68-043934.9	580
839	XMDS	J022315.3-044320	02 23 15.5 -04 43 18.4	48.2 10.8	22.3 9.8	3.9	9.0	B 0 2	2.1	25.45	0.16	24.24	0.36	1									839
310	XMDS	J022316.3-042007	02 23 16.4 -04 20 07.8	121.8 15.7	1.0 7.2	12.2	0.5	B 0 0	1.8												1	BD-04 388	310
846	XMDS	J022317.2-044035	02 23 17.4 -04 40 33.5	167.2 18.0	72.4 13.3	17.0	37.3	B 0 1	1.1	18.47	0.00	18.23	0.00	1									846
828	XMDS	J022318.8-044616	02 23 19.0 -04 46 13.9	166.7 17.9	53.0 11.5	11.0	17.4	B 0 0	0.2	21.16	0.01	20.77	0.01	0									828
321	XMDS	J022319.2-041648	02 23 19.4 -04 16 48.4	55.8 11.4	9.6 8.4	4.1	3.6	B 0 1	1.1	22.62	0.04	21.73	0.03	0									321
820	XMDS	J022319.4-044732	02 23 19.6 -04 47 30.4	280.6 22.5	125.6 16.0	17.4	38.7	B 0 0	0.7	19.14	0.01	18.49	0.00	0									820
779	XMDS	J022321.8-045740	02 23 22.0 -04 57 38.4	20.1 9.1	98.7 14.7	1.9	47.2	CD 0 0	1.0	20.35	0.01	19.47	0.01	1									779
333	XMDS	J022324.9-041452	02 23 25.1 -04 14 52.2	70.5 12.9	28.5 9.8	4.7	9.4	B 0 1	0.6	21.83	0.02	21.31	0.02	0									333
550	XMDS	J022326.0-043537	02 23 26.0 -04 35 37.1	115.3 15.3	31.3 10.6	17.7	25.1	B 1 1	1.0	23.16	0.04	22.33	0.04	0									550
315	XMDS	J022326.3-041838	02 23 26.5 -04 18 38.5	51.6 11.5	8.9 8.4	4.6	4.0	B 0 0	0.5	24.29	0.08	23.27	0.07	0									315
782	XMDS	J022326.3-045708	02 23 26.5 -04 57 05.1	122.8 15.8	62.2 12.3	11.4	29.1	B 0 0	1.5	20.74	0.01	20.49	0.01	1									782
789	XMDS	J022329.1-045452	02 23 29.3 -04 54 50.8	245.2 21.2	52.0 11.5	22.0	24.0	B 2 1	1.2	20.25	0.01	19.92	0.01	0									789
789									2	3.5	22.13	0.02	21.45	0.02	0								789
571	XMDS	J022330.2-043004	02 23 30.3 -04 30 04.2	121.7 15.7	47.0 11.7	14.0	27.6	B 1 0	1.7	20.18	0.01	19.94	0.01	0									571
825	XMDS	J022330.6-044633	02 23 30.8 -04 46 31.8	303.2 23.3	100.4 14.8	20.5	33.6	B 0 1	0.8	24.44	0.12	23.37	0.08	0									825
840	XMDS	J022330.9-044235	02 23 31.1 -04 42 33.0	59.1 12.4	37.6 10.6	6.3	20.8	B 2 2	5.4	24.77	0.14	24.59	0.24	0									840
780	XMDS	J022332.0-045740	02 23 32.2 -04 57 38.7	184.5 18.7	49.5 11.6	18.1	24.6	B 0 0	0.7	19.99	0.01	19.63	0.01	1									780
354	XMDS	J022332.2-041022	02 23 32.3 -04 10 22.1	128.4 16.2	9.3 7.6	7.7	2.8	B 0 0	0.3	17.77	0.00	16.93	0.00	2									354
807	XMDS	J022333.0-044924	02 23 33.2 -04 49 22.5	143.8 16.7	48.0 11.3	11.2	17.9	B 2 1	0.9	20.60	0.01	20.26	0.01	0									807
573	XMDS	J022336.2-042943	02 23 36.3 -04 29 43.8	50.4 11.2	4.1 7.1	6.4	2.7	B 2 1	1.8	16.06	0.00	16.89	0.00	2									573
316	XMDS	J022336.7-041828	02 23 36.9 -04 18 28.1	44.8 10.8	26.7 9.6	4.8	14.3	B 1 0	0.7	21.16	0.01	21.18	0.02	0									316
357	XMDS	J022337.3-040938	02 23 37.5 -04 09 38.2	100.5 14.7	2.0 7.1	6.5	0.6	B 0 0	1.5	15.40	0.00	14.67	0.00	2									357
849	XMDS	J022337.3-044008	02 23 37.6 -04 40 06.7	82.7 13.6	21.2 9.5	9.3	12.0	B 0 0	1.2	20.03	0.01	19.66	0.01	0									849
369	XMDS	J022337.7-040513	02 23 37.9 -04 05 13.1	97.8 14.3	25.4 9.4	9.8	13.4	B 2 1	0.6	24.62	0.14	24.98	0.44	0									369
574	XMDS	J022340.2-042851	02 23 40.3 -04 28 51.1	45.9 11.0	12.7 8.9	4.3	5.9	B 0 1	0.3	21.33	0.01	21.11	0.02	0									574
590	XMDS	J022340.3-042419	02 23 40.4 -04 24 19.1	98.1 14.3	25.4 10.0	11.0	14.6	B 0 0	0.5	24.37	0.10	23.07	0.07	0									590
596	XMDS	J022340.6-042255	02 23 40.7 -04 22 55.2	63.2 12.2	22.7 9.2	8.4	15.3	B 0 0	1.6	19.27	0.01	19.15	0.01	0									596
566	XMDS	J022341.8-043050	02 23 41.9 -04 30 50.7	104.7 14.9	17.1 9.1	9.5	7.8	B 0 2	2.7	21.10	0.01	20.37	0.01	0									566
566									2	3.6	21.24	0.01	20.43	0.01	0								566
566									2	1.2	18.88	0.00	18.09	0.00	0								566
551	XMDS	J022342.0-043533	02 23 42.1 -04 35 33.7	109.1 14.9	45.3 11.1	12.5	26.5	B 0 1	1.2	20.77	0.01	20.71	0.02	0									551
801	XMDS	J022344.4-045120	02 23 44.6 -04 51 18.2	114.6 15.3	45.5 11.2	8.8	17.4	B 0 2	1.1	21.97	0.02	21.50	0.02	0									801
801									2	4.4	18.69	0.00	17.57	0.00	0								801
558	XMDS	J022347.0-043346	02 23 47.0 -04 33 46.1	62.9 12.4	27.5 9.8	5.9	13.0	B 0 1	1.9	19.52	0.01	19.02	0.01	0									558
558									2	4.8	22.16	0.03	21.57	0.02	0								558
558									2	4.8	20.63	0.01	20.26	0.01	0								558
585	XMDS	J022348.9-044117	02 23 48.1 -04 41 17.8	45.6 10.9	4.8 7.8	11.1	5.8	B 2 2	2.6	24.71	0.14	24.36	0.18	0									585
585									2	2.6	24.49	0.12	23.52	0.09	0								585
579	XMDS	J022349.0-042658	02 23 49.0 -04 26 58.9	90.8 13.9	15.5 9.3	7.9	6.8	B 0 1	0.8	21.75	0.02	21.35	0.02	0									579
848	XMDS	J022349.0-044010	02 23 49.2 -04 40 08.1	62.1 12.4	20.3 9.9	8.7	14.7	B 2 2	4.7	21.27	0.02	20.45	0.01	0									848
578	XMDS	J022350.7-042703	02 23 50.7 -04 27 03.8	189.6 18.9	32.0 10.6	15.9	13.5	B 0 0	0.4	21.46	0.01	21.41	0.02	0									578
564	XMDS	J022350.7-043157	02 23 50.8 -04 31 57.9	162.9 17.8	52.7 11.7	13.4	21.8	B 0 1	0.2	19.36	0.01	19.23	0.01	0									564
786	XMDS	J022350.6-045528	02 23 50.9 -04 55 26.1	61.4 12.2	12.1 8.7	6.5	6.4	B 0 1	0.7	21.49	0.02	21.23	0.02	0									786
822	XMDS	J022350.9-044731	02 23 51.1 -04 47 29.8	87.5 13.8	9.5 8.1	7.7	4.2	B 0 0	0.1	20.42	0.01	20.10	0.01	0									822
314	XMDS	J022351.5-041840	02 23 51.7 -04 18 40.4	53.3 12.1	10.0 9.4	6.7	6.5	B 1 1	4.1	22.89	0.03	22.55	0.04	0									314
569	XMDS	J022352.1-043032	02 23 52.2 -04 30 32.9	72.4 13.3	20.9 9.2	7.8	10.8	B 2 0	1.5	19.65	0.01	19.34	0.01	0									569
788	XMDS	J022353.7-045510	02 23 53.9 -04 55 08.2	273.7 22.4	82.0 14.0	31.8	48.7	B 2 1	0.4	21.64	0.02	21.43	0.02	0									788
547	XMDS	J022353.9-043610	02 23 53.9 -04 36 10.8	63.3 12.6	6.4 7.9	7.7	4.1	B 2 0	0.9	21.95	0.03	21.72	0.03	0									547
817	XMDS	J022354.5-044815	02 23 54.8 -04 48 13.9	185.3 18.8	36.1 10.8	17.4	17.0	B 0 1	1.2	18.15	0.00	18.01	0.00	1									817
351	XMDS	J022356.5-041105	02 23 56.7 -04 11 05.8	96.2 14.6	32.5 10.8	9.6	16.7	B 1 0	0.5	21.81	0.02	21.54	0.02	0									351
565	XMDS	J022356.8-043115	02 23 56.8 -04 31 15.1	131.5 16.3	45.1 11.3	11.5	19.3	B 2 2	1.5	23.38	0.06	22.80	0.06	0									565
565									2	3.3	22.49	0.04	22.12	0.05	0								565
560	XMDS	J022403.7-043303	02 24 03.8 -04 33 03.3	69.8 12.8	31.1 10.4	5.7	12.7	B 0 0	1.6	15.14	0.00	14.84	0.00	2	V							*MRK 1036	560
800	XMDS	J022403.8-045120	02 24 04.1 -04 51 18.3	137.9 16.4	34.6 10.7	15.9	20.3	B 0 1	0.2	21.43	0.02	20.95	0.02	0									800
341	XMDS	J022403.9-041326	02 24 04.1 -04 13 26.5	84.7 15.6	12.8 9.8	10.5	8.3	B 1 2	13.6														341
327	XMDS	J022405.2-041611	02 24 05.3 -04 16 11.1	49.9 11.6	7.0 8.8	7.1	5.2	B 1 2	1.6	25.40	0.25	24.69	0.32	0									327
327									2	3.8	25.83	0.29	24.92	0.30	0								327
327									2	5.3	23.77	0.06	23.12	0.07	0								327
327									2	4.7	24.03	0.07	23.12	0.07	0								327
556	XMDS	J022409.7-043404	02 24 09.8 -04 34 04.5	59.8 12.2	20.4 9.2	5.3	9.1	B 0 0	0.4	21.32	0.02	21.21											

572	XMDS	J022430.9-042956	02	24	30.9	-04	29	56.6	89.8	13.9	18.5	9.5	11.1	11.3	B	2	1	0.6	21.23	0.02	21.11	0.01	0	572		
572																2	5.2	22.78	0.04	22.34	0.04	0	572			
567	XMDS	J022432.4-043036	02	24	32.5	-04	30	36.9	208.0	19.5	47.6	11.7	23.6	27.2	B	1	1	0.1	20.22	0.01	19.91	0.02	0	567		
267	XMDS	J022433.7-040902	02	24	33.7	-04	09	03.9	54.3	11.6	27.7	9.8	3.5	8.9	B	0	2	1.5	23.02	0.05	22.25	0.06	1	267		
267																2	2.2	22.61	0.03	22.29	0.04	1	267			
254	XMDS	J022433.9-042049	02	24	33.1	-04	20	50.1	68.7	12.7	12.1	9.0	8.5	7.6	B	0	1	0.6	24.09	0.07	22.96	0.06	0	254		
563	XMDS	J022434.3-043158	02	24	34.3	-04	31	58.7	69.0	12.4	8.9	8.4	8.3	5.4	B	0	1	1.4	19.67	0.01	19.50	0.01	0	563		
58	XMDS	J022436.2-042511	02	24	36.2	-04	25	10.8	66.9	12.8	36.4	11.3	8.3	23.2	B	0	0	0.5	23.74	0.06	23.31	0.08	0	V	VIRMOS 1.4GHz J022436-042511	58
242	XMDS	J022437.8-041520	02	24	37.8	-04	15	21.9	47.8	11.6	37.0	10.6	4.1	15.6	B	2	2	3.4	23.19	0.05	22.31	0.05	0	242		
242																2	5.5	22.66	0.03	21.77	0.03	0	242			
242																2	5.1	24.30	0.11	23.83	0.16	0	242			
283	XMDS	J022438.0-040530	02	24	38.1	-04	05	31.9	110.9	15.1	25.6	9.8	8.4	9.7	B	0	1	1.5	23.17	0.04	22.94	0.06	0	283		
577	XMDS	J022438.9-042705	02	24	38.9	-04	27	05.8	965.0	39.5	182.5	18.9	133.0	126.7	B	2	0	0.9	17.80	0.00	17.41	0.00	2	V	APMUks(BJ) B022208.33-044038.7	577
60	XMDS	J022439.6-042401	02	24	39.7	-04	24	01.1	231.0	20.6	75.1	13.5	28.1	47.1	B	0	0	0.2	19.76	0.01	19.27	0.01	0	V	VIRMOS 1.4GHz J022439-042401	60
14	XMDS	J022440.6-043657	02	24	40.7	-04	36	56.8	114.9	15.7	23.7	10.2	15.3	16.0	B	2	0	0.5	19.74	0.01	19.58	0.01	0	14		
262	XMDS	J022443.4-040953	02	24	43.5	-04	09	54.3	152.3	17.1	9.5	7.7	9.6	3.0	B	0	1	0.1	21.04	0.01	20.71	0.01	1	262		
542	XMDS	J022445.4-043656	02	24	45.4	-04	36	56.8	47.5	11.2	0.0	7.0	8.8	0.0	B	0	4							542		
258	XMDS	J022447.4-041049	02	24	47.5	-04	10	50.7	67.1	12.6	49.9	11.3	4.4	16.4	B	0	0	0.3	21.17	0.01	21.17	0.02	1	258		
232	XMDS	J022449.2-041800	02	24	49.2	-04	18	01.5	202.0	19.4	42.0	11.4	20.0	21.1	B	0	0	0.4	20.96	0.01	20.40	0.01	0	232		
270	XMDS	J022449.2-040841	02	24	49.3	-04	08	42.8	69.7	12.4	37.0	10.1	4.7	12.6	B	0	1	0.4	21.80	0.03	21.18	0.03	0	270		
36	XMDS	J022449.8-043026	02	24	49.9	-04	30	26.2	136.9	16.5	84.9	14.2	11.3	35.5	B	1	0	0.7	19.94	0.01	19.95	0.01	0	36		
57	XMDS	J022450.8-042506	02	24	50.9	-04	25	06.3	52.3	11.6	0.0	6.6	5.0	0.0	B	0	1	3.7	23.12	0.04	22.66	0.05	0	57		
45	XMDS	J022450.9-042903	02	24	50.1	-04	29	03.1	64.7	12.4	26.4	10.5	5.3	10.8	B	0	1	0.5	21.50	0.01	21.37	0.02	0	45		
45																2	2.8	23.29	0.04	23.10	0.06	0	V	L VIRMOS 1.4GHz J022451-042900	45	
253	XMDS	J022451.9-041209	02	24	52.0	-04	12	10.2	145.9	16.8	42.8	10.8	13.0	18.6	B	2	0	0.6	19.24	0.00	18.92	0.00	1	V	VIRMOS 1.4GHz J022452-041210	253
282	XMDS	J022452.1-040518	02	24	52.1	-04	05	20.1	535.7	30.2	108.5	15.1	44.3	45.3	B	0	1	1.0	19.91	0.01	19.60	0.01	0	282		
231	XMDS	J022453.1-041810	02	24	53.1	-04	18	11.1	69.4	12.9	16.6	9.0	7.5	9.3	B	1	2	4.4	21.84	0.02	21.20	0.02	0	231		
277	XMDS	J022454.7-040628	02	24	54.8	-04	06	29.9	105.3	14.7	25.7	9.9	8.4	10.4	B	0	1	1.1	21.37	0.01	21.02	0.02	0	277		
277																2	2.0	24.11	0.11	23.57	0.14	0	V	VIRMOS 1.4GHz J022454-040631	277	
30	XMDS	J022455.6-043201	02	24	55.6	-04	32	01.3	45.9	11.1	24.9	9.7	3.5	9.6	B	0	1	3.2	21.61	0.02	21.42	0.02	0	30		
30																2	4.0	23.86	0.09	23.68	0.13	0	30			
69	XMDS	J022456.0-041946	02	24	56.1	-04	19	45.7	80.7	13.5	24.9	10.4	10.5	16.8	B	0	1	0.5	22.43	0.02	22.03	0.03	0	69		
233	XMDS	J022456.0-041725	02	24	56.1	-04	17	26.7	100.8	14.6	34.7	11.1	10.3	18.1	B	0	2	14.8						2MASX J02245534-0417351	233	
73	XMDS	J022459.2-044151	02	24	59.3	-04	41	51.3	58.3	12.2	19.6	9.7	8.3	14.2	B	2	0	1.8	21.87	0.02	21.38	0.02	0	73		
249	XMDS	J022500.8-041313	02	25	00.9	-04	13	14.4	47.8	11.1	8.8	8.2	4.0	3.7	B	0	1	0.6	22.29	0.02	22.16	0.03	0	249		
272	XMDS	J022501.6-040752	02	25	01.7	-04	07	53.5	197.6	19.1	59.1	12.0	16.9	25.8	B	1	1	0.9	19.50	0.01	19.49	0.01	0	272		
2	XMDS	J022503.0-044025	02	25	03.0	-04	40	25.4	90.1	14.0	6.8	8.5	11.0	4.2	B	2	1	1.6	21.85	0.02	21.48	0.02	0	V	VIRMOS 1.4GHz J022503-044025	2
281	XMDS	J022503.2-040538	02	25	03.3	-04	05	39.5	402.4	26.4	103.2	15.0	40.4	52.9	B	0	1	0.8	20.96	0.01	20.62	0.01	0	281		
13	XMDS	J022504.5-043707	02	25	04.5	-04	37	07.4	79.7	13.3	43.9	11.5	8.1	23.0	B	0	2	0.4	23.41	0.07	22.70	0.06	0	13		
47	XMDS	J022506.2-042847	02	25	06.3	-04	28	47.4	178.1	18.6	14.2	9.2	11.2	4.5	B	0	0	1.3	17.62	0.00	17.21	0.00	2	47		
16	XMDS	J022506.4-043621	02	25	06.5	-04	36	20.9	164.1	18.1	33.5	11.0	13.3	13.7	B	0	0	0.6	21.45	0.02	20.77	0.01	0	16		
23	XMDS	J022507.1-043405	02	25	07.2	-04	34	05.2	83.5	13.8	25.3	10.1	5.7	8.8	B	0	0	2.6	21.58	0.02	20.94	0.01	0	23		
237	XMDS	J022507.9-041629	02	25	08.0	-04	16	30.1	60.6	12.1	16.5	8.8	7.0	9.8	B	0	2	0.7	24.51	0.10	22.95	0.07	1	237		
237																2	2.4	24.96	0.16	23.25	0.08	1	237			
271	XMDS	J022509.5-040836	02	25	09.6	-04	08	37.9	134.7	16.8	35.9	10.9	13.2	18.0	B	1	1	1.3	20.09	0.01	19.87	0.01	0	271		
18	XMDS	J022510.6-043549	02	25	10.7	-04	35	49.3	205.3	19.7	78.8	13.9	15.4	29.7	B	0	0	0.3		24.86	0.29	0	V	VIRMOS 1.4GHz J022510-043548	18	
40	XMDS	J022510.6-042928	02	25	10.7	-04	29	28.3	55.0	11.8	89.2	14.1	3.3	26.6	CD	0	2	1.9	23.85	0.08	23.25	0.10	0	40		
40																2	2.9	24.38	0.12	23.50	0.12	0	40			
71	XMDS	J022511.9-041911	02	25	11.1	-04	19	10.5	67.4	13.0	65.7	13.2	8.2	41.3	B	0	2	1.4	24.17	0.09	23.59	0.11	0	71		
71																2	4.1	22.06	0.02	21.75	0.03	0	71			
746	XMDS	J022512.6-044633	02	25	12.6	-04	46	33.8	36.5	10.5	102.9	15.3	5.2	74.8	CD	1	0	1.0	20.03	0.01	19.42	0.01	0	746		
742	XMDS	J022514.3-044659	02	25	14.3	-04	47	00.0	233.2	20.9	60.1	12.7	31.6	41.4	B	1	0	1.0	18.50	0.00	18.18	0.00	0	V	VIRMOS 1.4GHz J022514-044659	742
54	XMDS	J022514.3-042712	02	25	14.4	-04	27	11.6	193.8	19.1	8.0	8.1	12.1	2.5	B	0	0	1.7	16.05	0.00	17.42	0.01	2	54		
736	XMDS	J022515.0-044741	02	25	15.0	-04	47	41.8	85.5	13.9	21.1	9.6	11.2	14.0	B	1	1	1.1	21.47	0.02	20.96	0.02	0	736		
736																2	3.6	23.31	0.05	21.46	0.02	0	736			
3	XMDS	J022515.3-044008	02	25	15.4	-04	40	07.6	79.7	13.5	22.3	10.2	8.4	12.1	B	0	0	1.3	19.71	0.01	19.48	0.01	0	3		
286	XMDS	J022516.5-040511	02	25	16.6	-04	05	12.9	122.3	15.8	27.5	9.9	16.2	18.9	B	1	0	1.4	20.88	0.01	20.07	0.01	0	286		
44	XMDS	J022516.8-042910	02	25	16.9	-04	29	10.3	60.4	12.4	10.3	8.6	3.6	3.0	B	0	1	0.7	18.73	0.00	18.17	0.00	0	V	VIRMOS 1.4GHz J022516-042908	44
740	XMDS	J022519.3-044714	02	25	19.3	-04	47	14.8	95.5	14.4	27.4	10.0	11.9	17.3	B	0	0	1.4	20.90	0.01	20.50	0.01	0	740		
250	XMDS	J022520.7-041246	02	25	20.7	-04	12	47.3	136.6	16.7																

	XMDS	J022552.1-042249	02 25 52.2 -04 22 49.0	102.8 14.9	15.0 9.4	12.8	9.8	B 1 1	0.8 21.01 0.01 20.79 0.02 0		63
731	XMDS	J022554.1-044921	02 25 54.2 -04 49 21.9	49.6 11.8	34.9 10.3	3.7	12.5	B 2 2	0.7 22.44 0.03 21.73 0.03 0		731
731								2	1.8 23.16 0.04 22.81 0.05 0		731
7	XMDS	J022555.3-043920	02 25 55.4 -04 39 19.5	50.8 11.5	0.0 5.9	9.3	0.0	B 2 1	1.6 20.64 0.01 20.50 0.01 0		7
28	XMDS	J022555.9-043218	02 25 56.0 -04 32 18.3	110.9 15.2	21.8 10.3	11.6	11.9	B 1 0	2.0 20.42 0.01 20.13 0.01 0		28
738	XMDS	J022556.1-044724	02 25 56.1 -04 47 24.7	350.3 24.9	96.3 14.4	24.5	33.4	B 0 0	0.2 20.48 0.01 20.17 0.01 0	V	VIRMOS 1.4GHz J022556-044723 738
56	XMDS	J022556.3-042540	02 25 56.3 -04 25 39.7	98.8 14.8	20.9 10.3	11.5	12.7	B 1 2	1.5 21.50 0.01 21.08 0.01 0		56
56								2	2.4 23.27 0.04 22.26 0.03 0		56
758	XMDS	J022557.1-044300	02 25 57.1 -04 43 01.2	62.4 12.4	14.6 8.7	6.0	7.1	B 0 0	1.9 20.60 0.01 20.39 0.01 0		758
187	XMDS	J022557.4-041340	02 25 57.5 -04 13 39.6	62.2 11.9	7.6 6.5	14.9	8.6	B 2 0	1.1 20.32 0.01 19.83 0.01 0		187
730	XMDS	J022557.6-045005	02 25 57.6 -04 50 05.5	91.7 14.2	24.3 9.6	5.9	7.8	B 0 0	0.3 19.47 0.00 19.38 0.00 0	V	VIRMOS 1.4GHz J022557-045005 730
755	XMDS	J022600.1-044412	02 26 00.2 -04 44 13.3	215.0 19.9	60.8 12.1	19.6	28.1	B 2 2	2.4 25.33 0.27 24.44 0.21 0		755
755								2	2.7 23.03 0.03 22.04 0.02 0		755
27	XMDS	J022605.4-043226	02 26 05.5 -04 32 25.8	69.6 13.0	24.0 10.4	9.2	16.8	B 2 1	1.1 20.83 0.01 20.31 0.01 1		27
27								2	2.5 22.68 0.03 22.32 0.03 1		27
211	XMDS	J022606.6-040259	02 26 06.6 -04 02 58.5	55.1 11.3	3.6 6.3	12.9	4.2	B 2 2	3.7 18.84 0.00 18.66 0.00 0		211
709	XMDS	J022606.7-045722	02 26 06.7 -04 57 23.4	219.5 20.8	41.3 11.0	22.2	21.2	B 1 1	0.3 20.70 0.03 20.15 0.01 1		709
488	XMDS	J022606.9-043316	02 26 06.9 -04 33 17.1	59.1 11.6	8.5 7.7	12.6	9.2	B 1 2	5.7		BD-05 456 488
749	XMDS	J022606.9-044542	02 26 06.9 -04 45 43.0	50.5 11.4	27.8 9.3	4.0	11.1	B 0 2	0.7 23.77 0.08 22.81 0.05 0		749
749								2	2.6 24.49 0.14 24.15 0.14 0		749
749								2	2.7 25.00 0.18 24.67 0.21 0		749
749								2	2.3 24.01 0.11 23.32 0.09 0		749
185	XMDS	J022607.4-041405	02 26 07.4 -04 14 05.1	71.5 12.3	26.5 9.0	12.5	23.0	B 0 1	0.4 23.85 0.07 23.26 0.08 0		185
179	XMDS	J022607.7-041843	02 26 07.7 -04 18 42.9	328.6 23.8	90.1 13.6	87.3	120.8	B 0 0	0.4 19.47 0.01 18.96 0.01 0		179
748	XMDS	J022610.9-044550	02 26 10.1 -04 45 51.1	101.6 14.6	49.9 11.3	8.3	20.5	B 0 1	0.7 22.36 0.04 22.15 0.04 0		748
485	XMDS	J022612.6-043401	02 26 12.7 -04 34 02.0	46.9 10.7	8.8 7.6	9.2	8.7	B 0 1	1.0 20.57 0.01 20.36 0.01 0		485
718	XMDS	J022615.1-045355	02 26 15.1 -04 53 55.7	100.5 14.8	51.3 11.5	8.4	21.5	B 0 1	0.3 21.92 0.03 21.13 0.02 0	V	VIRMOS 1.4GHz J022615-045353 718
739	XMDS	J022617.1-044724	02 26 17.2 -04 47 24.9	25.5 9.1	56.2 11.7	2.1	23.4	CD 0 0	0.9 17.85 0.00 17.39 0.00 2	V	APMUKS (BJ) B022346.77

L.Chiappetti et al.: The XMDS/VVDS 4σ catalogue, *Online Material p 5*

102	XMDS	J022702.7-041511	02 27 02.9	-04 15 11.3	114.2	15.7	25.2	10.0	9.3	10.5	B 2 2	4.6	23.45	0.05	23.43	0.08	0	102
102											2	6.0	25.39	0.21	25.30	0.39	0	102
102											2	4.1	22.76	0.03	22.22	0.03	0	102
89	XMDS	J022703.9-041709	02 27 04.1	-04 17 09.9	60.0	12.1	6.6	8.2	5.3	3.0	B 2 0	0.2	21.61	0.02	21.35	0.02	0	89
117	XMDS	J022704.0-042035	02 27 04.2	-04 20 35.1	73.8	13.3	5.5	8.4	8.2	3.2	B 1 1	1.8	25.79	0.30	25.11	0.34	0	117
81	XMDS	J022706.3-041924	02 27 06.5	-04 19 24.6	45.9	11.3	12.3	8.6	4.4	6.0	B 0 0	0.6	20.99	0.01	21.09	0.02	0	81
149	XMDS	J022707.2-040438	02 27 07.4	-04 04 38.6	179.8	18.5	54.5	12.3	12.8	19.6	B 0 0	1.3	20.29	0.01	19.70	0.01	0	149
107	XMDS	J022708.6-041359	02 27 08.7	-04 13 59.9	52.4	11.6	17.6	9.3	3.3	5.6	B 0 0	1.6	22.37	0.03	21.67	0.02	0	107
489	XMDS	J022708.9-043245	02 27 08.9	-04 32 45.9	51.9	11.4	15.7	8.4	9.1	13.7	B 0 1	1.1	21.45	0.02	20.91	0.02	0	489
91	XMDS	J022710.0-041649	02 27 10.2	-04 16 49.5	138.2	16.6	51.6	11.7	11.6	22.2	B 2 2	2.1	20.35	0.01	19.87	0.01	0	91
91											2	1.0	22.51	0.03	21.76	0.02	0	91
483	XMDS	J022710.4-043416	02 27 10.5	-04 34 17.4	97.1	14.2	0.0	5.8	19.1	0.0	B 0 2	2.3	26.07	0.45	25.74	0.49	0	483
483											2	4.6	22.50	0.03	22.03	0.03	0	483
1204	XMDS	J022710.5-045531	02 27 10.6	-04 55 31.9	55.8	11.6	8.5	7.9	4.1	3.1	B 0 0	2.7	18.01	0.00	16.58	0.00	2	1204
1226	XMDS	J022711.7-045038	02 27 11.8	-04 50 38.6	916.8	38.6	289.9	22.8	54.5	85.5	B 0 1	1.1	20.53	0.01	20.42	0.01	0	1226
1265	XMDS	J022712.6-044221	02 27 12.7	-04 42 21.7	115.4	15.4	138.6	16.8	10.4	62.8	CD 0 0	0.9	18.40	0.00	17.91	0.00	0	V VIRMOS 1.4GHz J022712-044221 1265
1246	XMDS	J022712.8-044636	02 27 12.9	-04 46 37.1	285.8	22.8	73.2	13.1	18.4	23.4	B 0 0	1.6	17.99	0.00	17.69	0.00	0	V L NVSS J022713-044637 1246
133	XMDS	J022713.1-040912	02 27 13.2	-04 09 12.8	214.7	19.9	84.6	14.0	11.6	22.9	B 0 2	2.5	23.58	0.06	23.13	0.07	0	133
133											1	0.7	23.21	0.05	22.35	0.04	0	133
1276	XMDS	J022713.6-043910	02 27 13.7	-04 39 10.9	113.6	15.4	5.7	7.3	13.7	3.5	B 0 0	0.7						BD-05 461 1276
444	XMDS	J022714.2-042645	02 27 14.2	-04 26 45.7	220.3	20.5	19.0	9.4	26.6	11.7	B 1 2							444
1252	XMDS	J022716.0-044539	02 27 16.0	-04 45 39.6	552.8	30.6	137.2	16.8	37.4	46.2	B 0 0	1.6	20.37	0.01	20.02	0.01	0	1252
417	XMDS	J022716.2-043244	02 27 16.3	-04 32 44.2	44.3	11.0	16.6	9.3	5.2	10.0	B 1 2	3.4	20.29	0.01	19.87	0.01	0	417
425	XMDS	J022718.9-043132	02 27 18.1	-04 31 32.1	90.6	14.2	25.0	9.9	9.5	13.3	B 1 2	2.0	21.00	0.01	20.74	0.01	0	425
425											2	2.4	24.12	0.10	23.56	0.09	0	425
425											2	3.5	22.39	0.04	21.67	0.02	0	425
1253	XMDS	J022720.6-044537	02 27 20.7	-04 45 37.9	104.0	14.8	3.4	7.3	7.1	1.2	B 0 0	1.4	15.52	0.00	15.16	0.00	2	V 2MASX J02272073-0445374 1253
1268	XMDS	J022721.7-044153	02 27 21.8	-04 41 54.0	65.0	12.1	16.8	8.5	6.1	8.0	B 0 1	0.8						V VIRMOS 1.4GHz J022721-044153 1268
1201	XMDS	J022723.4-045608	02 27 23.5	-04 56 08.6	112.5	15.3	44.3	11.3	8.4	16.6	B 0 1	1.6	21.57	0.02	21.34	0.02	0	1201
1201											2	5.0	20.11	0.01	19.51	0.01	0	1201
1216	XMDS	J022723.6-045216	02 27 23.6	-04 52 17.0	111.8	15.4	25.3	10.2	6.8	7.6	B 0 1	1.8	21.49	0.02	21.19	0.02	0	1216
155	XMDS	J022723.6-040214	02 27 23.7	-04 02 14.7	107.6	14.9	13.2	8.5	9.1	5.7	B 0 0	1.1	19.77	0.01	19.54	0.01	1	V VIRMOS 1.4GHz J022723-040213 155
412	XMDS	J022724.0-043356	02 27 24.1	-04 33 56.1	47.1	11.6	8.5	8.4	4.7	4.3	B 1 0	0.4	23.85	0.08	23.38	0.06	0	412
476	XMDS	J022725.1-043656	02 27 25.2	-04 36 57.4	51.4	11.2	4.4	7.1	14.9	6.4	B 0 4							476
1248	XMDS	J022725.4-044619	02 27 25.5	-04 46 19.7	466.3	28.1	137.9	16.8	31.4	46.2	B 0 0	0.2	15.76	0.00	15.30	0.00	2	2MASX J02272548-0446194 1248
1260	XMDS	J022725.9-044424	02 27 26.0	-04 44 25.2	132.0	16.3	25.9	9.4	10.0	9.9	B 0 0	0.2	23.41	0.05	22.22	0.03	0	1260
414	XMDS	J022726.3-043327	02 27 26.4	-04 33 27.6	88.4	14.3	34.3	10.9	8.3	16.3	B 1 1	0.9	19.58	0.01	19.04	0.00	0	414
445	XMDS	J022727.7-042655	02 27 27.8	-04 26 55.7	63.1	12.9	15.9	9.2	7.8	9.5	B 2 1	1.2	20.95	0.01	20.72	0.01	0	445
445											2	5.6	20.61	0.01	19.78	0.01	0	445
139	XMDS	J022727.7-040806	02 27 27.8	-04 08 06.3	335.4	24.1	73.7	13.0	20.9	22.9	B 0 1	0.7	20.57	0.01	19.96	0.01	0	139
420	XMDS	J022729.2-043225	02 27 29.2	-04 32 25.1	156.7	18.2	41.0	11.4	14.6	19.2	B 2 1	1.5	19.12	0.01	19.04	0.00	0	420
455	XMDS	J022730.9-042503	02 27 30.1	-04 25 03.8	74.3	13.2	0.0	7.4	7.1	0.0	B 0 2	5.5	24.36	0.09	23.73	0.10	0	455
455											2	5.8	25.26	0.19	25.38	0.44	0	455
1231	XMDS	J022731.9-044957	02 27 31.9	-04 49 57.6	108.0	15.2	31.7	9.9	6.7	9.9	B 0 0	0.8	21.79	0.02	20.95	0.01	0	V VIRMOS 1.4GHz J022731-044958 1231
406	XMDS	J022736.7-043544	02 27 32.7	-04 35 44.7	56.3	12.9	86.4	14.2	5.2	40.4	CD 0 1	1.5	22.30	0.03	21.91	0.02	0	V VIRMOS 1.4GHz J022732-043544 406
409	XMDS	J022733.8-043422	02 27 33.8	-04 34 22.2	49.0	11.4	11.1	8.3	4.3	4.9	B 1 0	1.2	21.48	0.02	20.67	0.01	0	409
454	XMDS	J022733.9-042521	02 27 33.1	-04 25 21.7	183.6	18.8	24.7	10.0	16.0	10.9	B 0 0	1.7	20.38	0.01	20.23	0.01	0	454
463	XMDS	J022734.2-042226	02 27 34.2	-04 22 26.7	57.9	12.2	14.5	9.4	6.1	7.8	B 0 1	1.8	20.58	0.01	20.32	0.01	0	463
128	XMDS	J022734.1-041021	02 27 34.3	-04 10 21.3	57.1	12.2	20.8	9.5	3.4	6.3	B 0 2	1.6	24.02	0.09	23.56	0.10	0	128
128											2	1.8	24.71	0.12	24.24	0.14	0	128
111	XMDS	J022735.6-041317	02 27 35.7	-04 13 17.0	107.1	15.1	99.4	14.7	7.0	33.0	B 0 2	0.8	24.62	0.16	24.47	0.26	0	V VIRMOS 1.4GHz J022735-041317 111
111											2	0.4	24.86	0.18	24.47	0.23	0	V VIRMOS 1.4GHz J022735-041317 111
111											2	2.3	25.25	0.23	24.91	0.30	0	V VIRMOS 1.4GHz J022735-041317 111
111											2	2.6	24.84	0.16	24.50	0.20	0	V VIRMOS 1.4GHz J022735-041317 111
85	XMDS	J022735.6-041828	02 27 35.7	-04 18 28.6	83.5	13.6	23.3	10.0	7.8	11.2	B 0 1	1.1	19.78	0.01	19.44	0.01	0	85
80	XMDS	J022735.6-042023	02 27 35.8	-04 20 23.0	84.6	13.9	31.8	10.8	9.3	18.1	B 0 1	0.3	23.04	0.04	22.02	0.03	0	80
120	XMDS	J022735.7-041122	02 27 35.8	-04 11 22.6	889.5	38.1	288.0	22.8	55.2	90.5	B 0 2	1.7	21.60	0.02	21.33	0.02	0	V L NVSS J022736-041117 120
120											2	0.6	23.96	0.09	23.26	0.08	0	V L NVSS J022736-041117 120
448	XMDS	J022736.8-042633	02 27 36.9	-04 26 33.0	55.3	12.1	7.4	8.2	4.3	2.9	B 0 0	2.0	20.85	0.01	20.66	0.01	0	448
1227	XMDS	J022736.8-045033	02 27 36.9	-04 50 34.2	72.7	13.6	71.3	13.3	4.9	24.1	B 0 0	1.0	20.04	0.01	19.44	0.01	0	V VIRMOS 1.4GHz J022736-045032 1227
430	XMDS	J022737.1-043031	02 27 37.1	-04 30 31.5	119.7	15.7	41.1	10.8	11.9	20.5	B 2 0	1.1	22.76	0.03	22.10	0.02	0	430
116	XMDS	J022737.0-041218	02 27 37.1	-04 12 18.6	45.7	11.3	18.6	8.8	3.0	6.1	B 0 0	0.4	23.41	0.05	23.04	0.05	0	116
96	XMDS	J022738.2-041617	02 27 38.3	-04 16 17.8	45.0	10.9	0.0	6.4	3.7	0.0	B 0 0	2.4	14.90	0.00	16.29	0.00	2	96
131	XMDS	J022738.8-040941	02 27 38.1	-04 09 41.1	93.9	14.3	21.7	9.8	6.1	7.1	B 0 0	0.7	21.54	0.02	21.51	0.02	0	131
1217	XMDS	J022740.2-045129	02 27 40.3	-04 51 29.7	462.2	29.5	23.8	10.9	33.5	8.7	B 0 1	1.1	17.97	0.00	17.35	0.00	2	V APMUKS(BJ) B022510.04-050456.5 1217
83	XMDS	J022740.4-041858	02 27 40.6	-04 18 58.5	76.0	13.1	25.2	10.2	7.8	13.4	B 0 2	0.7	24.25	0.09	23.69	0.11	0	83
83											2	2.6	22.45	0.03	22.16	0.03	0	83
154	XMDS	J022740.4-040250	02 27 40.6	-04 02 50.9	139.4	16.5	15.1	9.0	15.4	8.7	B 2 0	0.5	18.28	0.00	18.11	0.00	0	V NVSS J022740-040250 154
403	XMDS	J022742.1-043607	02 27 42.1	-04 36 07.7	68.3	12.7	45.2	11.9	5.7	19.2</								

L.Chiappetti et al.: The XMDS/VVDS 4 σ catalogue, *Online Material p 6*

397	XMDS	J022755.8-043800	02	27	55.8	-04	38	00.8	93.9	14.2	17.3	8.7	8.2	7.7	B	0	2	1.4	25.28	0.22	25.50	0.34	0	397	
397																	2	1.7	24.42	0.14	24.46	0.18	0	397	
397																	2	2.1	25.23	0.21	24.92	0.20	0	397	
397																	2	3.3	25.40	0.26	25.05	0.24	0	397	
1209	XMDS	J022756.0-045436	02	27	56.1	-04	54	36.6	58.2	12.0	16.6	9.3	6.6	9.7	B	1	0	1.3	21.22	0.02	21.31	0.02	0	1209	
136	XMDS	J022758.4-040851	02	27	58.6	-04	08	51.6	50.2	11.2	10.6	9.0	5.1	5.6	B	1	2	2.1	21.39	0.01	21.08	0.01	0	136	
136																	2	2.7	22.93	0.04	22.53	0.04	0	136	
427	XMDS	J022758.6-043112	02	27	58.6	-04	31	12.0	161.5	18.0	46.8	11.7	9.2	13.4	B	0	1	0.2	24.37	0.10	23.30	0.05	0	427	
442	XMDS	J022758.8-042754	02	27	58.8	-04	27	54.7	63.2	12.7	23.8	9.5	3.8	7.1	B	0	0	0.8	25.99	0.38	25.09	0.34	0	V	VIRMOS 1.4GHz J022758-042754 442
137	XMDS	J022802.0-040809	02	28	02.1	-04	08	09.1	99.8	14.6	25.8	10.1	11.2	15.2	B	1	1	1.1	20.90	0.01	20.56	0.01	0	137	
453	XMDS	J022802.3-042546	02	28	02.4	-04	25	46.8	29.9	9.7	58.0	11.9	2.1	20.7	CD	0	0	1.6	21.32	0.02	20.71	0.01	0	453	
470	XMDS	J022804.5-041818	02	28	04.6	-04	18	18.1	114.6	15.8	35.7	11.3	15.3	24.9	B	0	2	1.3	25.14	0.22	24.53	0.25	0	470	
470																	2	3.7	23.19	0.04	22.96	0.06	0	470	
470																	2	0.2	25.66	0.28	25.00	0.31	0	470	
394	XMDS	J022805.8-043924	02	28	05.9	-04	39	24.1	72.4	13.3	15.2	9.1	7.3	7.8	B	0	2	4.7	26.43	0.64	25.63	0.38	0	394	
394																	2	5.5	25.59	0.31	24.51	0.15	0	394	
394																	2	5.8	25.68	0.33	24.99	0.23	0	394	
112	XMDS	J022809.0-041232	02	28	09.1	-04	12	32.9	533.6	30.2	157.6	18.2	125.9	194.5	B	2	1	1.9	19.24	0.01	19.05	0.01	0	112	
101	XMDS	J022809.4-041524	02	28	09.5	-04	15	24.1	124.3	16.2	36.4	11.9	18.3	28.5	B	1	0	1.0	22.90	0.03	21.74	0.02	0	101	
416	XMDS	J022812.2-043230	02	28	12.3	-04	32	30.8	266.5	22.1	88.0	13.8	17.5	29.2	B	0	1	3.3	20.96	0.01	20.49	0.01	0	416	
914	XMDS	J022812.5-045717	02	28	12.5	-04	57	17.1	91.6	14.1	27.0	10.0	17.7	25.9	B	2	2	1.0	26.21	0.54	25.14	0.29	1	914	
914																	2	3.3	21.95	0.02	21.49	0.02	1	914	
467	XMDS	J022815.1-041942	02	28	15.1	-04	19	42.5	65.5	12.4	6.5	7.6	9.8	4.9	B	2	1	0.3	19.72	0.01	19.73	0.01	0	467	
449	XMDS	J022815.2-042617	02	28	15.3	-04	26	17.0	55.9	12.0	44.6	11.0	4.2	16.8	B	0	2	1.9	24.67	0.12	23.94	0.12	0	449	
449																	2	3.7	24.11	0.15	23.59	0.21	0	449	
472	XMDS	J022824.5-041810	02	28	24.5	-04	18	10.1	68.4	13.2	15.2	9.9	11.1	13.0	B	1	1	1.4	21.00	0.01	20.14	0.01	0	472	
395	XMDS	J022832.6-043825	02	28	32.7	-04	38	25.1	55.1	11.9	13.8	8.7	7.2	9.4	B	0	1	0.4	24.28	0.12	23.20	0.06	0	395	
410	XMDS	J022835.6-043414	02	28	35.6	-04	34	14.5	58.6	12.2	11.1	9.0	6.5	6.3	B	1	0	1.0	21.42	0.02	20.79	0.01	0	410	